arXiv:0904.0239v2 [math.GT] 18 Jul 2009

Brane Topological Field Theories and Hurwitz numbers for CW-complexes

Sergey M. Natanzon[1]

A.N.Belozersky Institute, Moscow State University
Independent University of Moscow
Institute Theoretical and Experimental Physics
natanzon@mccme.ru

1. Introduction

We expand Topological Field Theory of [5],[12] on some special CW-complexes (brane complexes). This Brane Topological Field Theory extends the Open Topological Field Theory from [9],[10],[1]. For arbitrary two-dimension brane complexes the Brane Topological Field Theory is the "open part" of Cyclic Foam Topological Field Theory from [11]. We prove that the Brane Topological Field Theory one-to-one corresponds to infinite dimensional Frobenius Algebras, graduated by CW-complexes of lesser dimension.

We define general and regular Hurwitz numbers of brane complexes and prove that they generate Brane Topological Field Theories. For general Hurwitz numbers corresponding algebra is an algebra of coverings of lesser dimension. For regular Hurwitz numbers the Frobenius algebra is an algebra of families of subgroups of finite groups.

2. Brane Topological Field Theory

2.1. **Simple CW-complexes.** A finite compact CW-complex is said to be *simple* if

- it is the closure of theirs cells of maximal dimension;
- the closure of any cell is a closed ball.

An isomorphism between simple CW-complexes is a homeomorphism conserving theirs cell structures. Zero-dimensional cells are called *vertexes*. Denote by Ω_b the set of all vertexes of Ω.

Denote by $|M|$ the cardinality of a set M. Following by Smith and Dold (see also [6]) we define *d-brunched covering* as a continuous map $h : X \to Y$ between two Hausdorff spaces and a continuous map $t : Y \to \mathrm{Sym}^d(X)$ such that: 1) $x \in th(x)$ for all $x \in X$, 2) $\mathrm{Sym}^d(h)(ty) = dy$ for all $y \in Y$. We say that a d-brunched covering is *finite* if Y assume a CW-complex structure such that the function $|h^{-1}(y)|$ is constant on any cell.

Lemma 2.1. *Any simple CW-complexes has a finite brunched covering over a ball.*

Proof. Let Ω be a simple CW-complex with vertexes $\{q_k|k \in K\}$. Consider a ball B and k different points $\{b_k \in \partial B|k \in K\}$. Then there exists a finite number isomorphic classes of simple CW-complexes with vertexes $\{b_k|k \in K\}$ that form a stratification of B. Consider a set $\{B_i|i \in I\}$ of representatives of theirs isomorphic classes.

Let $\{\Omega_j|j \in J\}$ be the set of closing of $\dim\Omega$-dimensional cells $\omega_j \subset \Omega$. Then for any $j \in J$ there exists an isomorphism of simple CW-complexes $h_j : \Omega_j \to B_{i(j)}$. Gluing now the manifolds $\{\Omega_j|j \in J\}$ along its boundary by maps $h_k^{-1}h_j$, using the same rules that

---

[1] Supported by grants RFBR-08-01-00110, NWO 047.011.2004.026 (05-02-89000-NWO-a), NSh-709.2008.1

manifolds $\{\Omega_j | j \in J\}$ form Ω. Then we get a simple CW-complex $\widetilde{\Omega}$ that isomorphic to Ω. The maps h_j generate a finite branching covering $h : \widetilde{\Omega} \to B$.

□

Theorem 2.1. *A Hausdorff space Ω has a stratification, generating a simple CW-complex if and only if there exists its finite branching covering over a ball $h : \Omega \to B$.*

Proof. Let $h : \Omega \to B$ be a finite branching covering. Construct a stratification of B onto cells, such that, the functions $|h^{-1}(y)|$ is constant on any cell. Preimages of these cells form a structure of simple CW-complex on Ω. The inverse affirmation follows from lemma 2.1.

□

2.2. **Brane complexes.** A simple CW-complex is said to be *oriented* if all cells of positive dimensional are oriented. Fix a set S of *colors*. An oriented complex Ω is called *colored complex* if a color $s(l) \in S$ is assigned to any its cell l of positive dimension, and for any connected component of Ω any two celles of the same dimension have different colors. *An isomorphism* between colored complexes is an isomorphism between simple CW-complexes, conserving the orientations and the colors of cells.

A system of cyclic orders on vertexes of connected components of a oriented complex Ω is called a *cyclic order on* Ω. A colored complex with a cyclic order is called *cyclic complex.*

A connected subcomplex $\gamma \subset \Omega$ of codimension 1 into cyclic complex Ω is called a *cut* if:

- the restriction of γ to any closed cell of $\omega \in \Omega$ either is empty, or homeomorphic to closed ball of codimension 1 in ω;
- γ divides a connected component of Ω into two connected components that splits its vertices into two nonempty groups, compatible with the cyclic order on Ω.

A cyclic complex is called a *brane complex*, if for any compatible with the cyclic order division of vertexes Ω there exists a cut that realize it.

The structure of CW-complex on Ω forms a structure of CW-complex on any its cut. A cut σ_q, dividing a vertex q from other vertexes, is called *vertex complex.* It separates a neighborhood U_q of q, that is a cone over σ_q. Any cell $\sigma \in \sigma_q$ divides an oriented cell $\omega \in \Omega$. Equip σ by orientation, as a boundary of connected component of $\omega \setminus \sigma$ that don't contain q. The colore structure on Ω generate a colore structure on σ_q. Thus, a brane complex generates a structure of connected colored complexes on its vertex complexes.

2.3. **Topological Field Theory.** Below we assume that all vector spaces are defined over a field $\mathbb{K} \supset \mathbb{Q}$. Let $\{X_m | m \in M\}$ be a finite set of $n = |M|$ vector spaces X_m over the field of complex numbers $\mathbb{C}$. The action of the symmetric group S_n on $\{1, \ldots, n\}$ induces its action on the sum of the vector spaces $\left(\oplus_\sigma X_{\sigma(1)} \otimes \cdots \otimes X_{\sigma(n)}\right)$, where σ runs over the bijections $\{1, \ldots, n\} \to M$, an element $s \in S_n$ takes $X_{\sigma(1)} \otimes \cdots \otimes X_{\sigma(n)}$ to $X_{\sigma(s(1))} \otimes \cdots \otimes X_{\sigma(s(n))}$. Denote by $\otimes_{m \in M} X_m$ the subspace of all invariants of this action. The vector space $\otimes_{m \in M} X_m$ is canonically isomorphic to the tensor product of all X_m in any fixed order; the isomorphism is the projection of $\otimes_{m \in M} X_m$ to the summand that is equal to the tensor product of X_m in that order.

Denote by $\Sigma = \Sigma(D, S)$ the set of all isomorphism classes of D-dimensional connected colored complexes. The inversion of the orientations generates the involution $* : \Sigma \to \Sigma$. Denote it by $\sigma \mapsto \sigma^*$.

Consider a family of finite-dimensional vector spaces $\{B_\sigma | \sigma \in \Sigma\}$ and a family of tensors $\{K^{\otimes}_\sigma \in B_\sigma \otimes B_{\sigma^*} | \sigma \in \Sigma\}$. Using these data, we define now a functor $\mathcal{V}$ from the monoidal category $\mathcal{S}$ of $(D+1)$-dimensional brane complexes to the category of vector spaces. This functor assigns the vector space $V_\Omega = (\otimes_{q \in \Omega_b} B_q)$ to any $(D+1)$-dimensional brane complex Ω. Here B_q is the copy of B_{σ_q} that is a vector space with a fixed isomorphism $B_q \to B_{\sigma_q}$.

We are going to describe all morphisms of the monoidal category $\mathcal{S}$ and morphisms of the category of vector spaces that correspond to it.

(1) *Isomorphism.* Define an isomorphism of brane complexes as an isomorphism $\phi : \Omega \to \Omega'$ of colored complexes, preserving the cyclic orders. Define $\mathcal{V}(\phi) = \phi_* : V_\Omega \to V_{\Omega'}$ as linear operator generated by the bijections $\phi|_{\Omega_b} : \Omega_b \to \Omega'_b$.

(2)*Cut.* Let Ω be a connected brane complex and $\gamma \subset \Omega$ be a cut. The cut γ is represented by two complexes γ_+ and γ_- on the closure $\overline{\Omega \setminus \gamma}$ of $\Omega \setminus \gamma$. Contract these complexes to points $q_+ = q_+[\gamma]$ and $q_- = q_-[\gamma]$, respectively. The contraction produces a brane complex $\Omega' = \Omega[\gamma]$. Its vertices $\Omega' = \Omega[\gamma]$ are the vertices of Ω and the points q_+, q_-. The cyclic order, orientation and the coloring of Ω induce an orientation and a coloring of Ω'. Thus, we can assume that Ω' is a brane complex and $V_{\Omega'} = V_\Omega \otimes B_{q_+} \otimes B_{q_-}$. The functor takes the morphism $\mathcal{V}(\eta)(x) = \eta_*(x) = x \otimes K^{\otimes}_\sigma$, where $\sigma = \sigma_{q_+} = \sigma^*_{q_-}$, to the morphism $\eta : \Omega \to \Omega'$.

(3) The tensor product in $\mathcal{S}$ defined by the disjoint union of brane complexes $\Omega' \otimes \Omega'' \to \Omega' \coprod \Omega''$ induces the tensor product of vector spaces $\theta_* : V_{\Omega'} \otimes V_{\Omega''} \to V_{\Omega' \sqcup \Omega''}$.

The functorial properties of $\mathcal{V}$ can be easily verified.

Fix a tuple of vector spaces and vectors $\{B_\sigma, K^{\otimes}_\sigma \in B_\sigma \otimes B_{\sigma^*} | \sigma \in \Sigma\}$, defining the functor $\mathcal{V}$. A family of linear forms $\mathcal{F} = \{\Phi_\Omega : V_\Omega \to \mathbb{K}\}$ defined for all brane complexes $\Omega \in \mathcal{S}$ is called a *Brane Topological Field Theory* if it satisfies the following axioms:

1° *Topological invariance.*

$$\Phi_{\Omega'}(\phi_*(x)) = \Phi_\Omega(x)$$

for any isomorphism $\phi : \Omega \to \Omega'$ of brane complexes.

2° *Non-degeneracy.*

Let Ω be a brane complex with only two vertices q_1, q_2. Then $\sigma_{q_2} = \sigma^*$ if $\sigma_{q_1} = \sigma$. Denote by $(.,.)_\sigma$ the bilinear form $(.,.)_\sigma : B_\sigma \times B_{\sigma^*} \to \mathbb{K}$, where $(x', x'')_\sigma = \Phi_\Omega(x'_{q_1} \otimes x''_{q_2})$. Axiom 2° asserts that the forms $(.,.)_\sigma$ are non-degenerated for all $\sigma \in \Sigma$.

3° *Cut invariance.*

$$\Phi_{\Omega'}(\eta_*(x)) = \Phi_\Omega(x)$$

for any cut morphism $\eta : \Omega \to \Omega'$ of brane complexes.

4° *Multiplicativity.*

$$\Phi_\Omega(\theta_*(x' \otimes x')) = \Phi_{\Omega'}(x')\Phi_{\Omega''}(x'')$$

for $\Omega = \Omega' \coprod \Omega''$, $x' \in V_{\Omega'}$, $x'' \in V_{\Omega''}$.

Note that a Topological Field Theory defines the tensors $\{K^{\otimes}_{\sigma} \in B_{\sigma} \otimes B_{\sigma^*} | \sigma \in \Sigma\}$, since it is not difficult to prove:

Lemma 2.2. *Let* $\{\Phi_{\Omega}\}$ *be a Brane Topological Field Theory. Then* $(K^{\otimes}_{\sigma}, x_1 \otimes x_2)_{\sigma} = (x_1, x_2)_{\sigma}$*, for all* $x_1 \in B_{\sigma}$*,* $x_2 \in B_{\sigma^*}$*.*

Remark 2.1. *Brane complexes with single cell of higher dimensional form subcategory into the subcategory (ball category) of brane complexes. Thus, we can consider a restriction Brane Topological Field Theory on ball category.*

3. Colored Frobenius Algebras

3.1. **Algebra.** We say that a connected brane complex Ω is a *compatible complex* for connected colored complexes $\sigma_1, \sigma_2, ..., \sigma_n$ if $\{\sigma_i\}$ are vertex of Ω and the numeration of $\{\sigma_i\}$ generates the cyclic order on Ω. Denote by $\Omega(\sigma_1, \sigma_2, ..., \sigma_n)$ the set of all isomorphism classes of compatible complexes for $\sigma_1, \sigma_2, ..., \sigma_n$. Then $\Omega(\sigma_1, \sigma_2, ..., \sigma_n)$ is either empty or consists of a single element.

Let $\Omega(\sigma_1, \sigma_2, \sigma_3, \sigma_4) \neq \emptyset$. Then there exist unique classes of cuts $\sigma_{(1,2|3,4)}, \sigma_{(4,1|2,3)} \in \Sigma$ such that $\Omega(\sigma_1, \sigma_2, \sigma_{(1,2|3,4)}) \neq \emptyset$, $\Omega(\sigma_{(3,4|1,2)}, \sigma_3, \sigma_4) \neq \emptyset$, $\Omega(\sigma_4, \sigma_1, \sigma_{(4,1|2,3)}) \neq \emptyset$, $\Omega(\sigma_{(4,1|2,3)}, \sigma_2, \sigma_3) \neq \emptyset$ and $\sigma_{(3,4|1,2)} = \sigma^*_{(1,2|3,4)}$, $\sigma_{(2,3|4,1)} = \sigma^*_{(4,1|2,3)}$.

Consider a tuple of finite dimensional vector spaces $\{B_{\sigma} | \sigma \in \Sigma\}$. Its direct sum $B_{\star} = \bigoplus_{\sigma \in \Sigma} B_{\sigma}$ is called a *colored vector space.* A colored vector space with a bilinear form $(.,.) : B_{\star} \times B_{\star} \to \mathbb{K}$ and a tree-linear form $(.,.,.) : B_{\star} \times B_{\star} \times B_{\star} \to \mathbb{K}$ is called a *Colored Frobenius Algebra* if

- the form $(.,.)$ is not-degenerate;
- $(B_{\sigma_1}, B_{\sigma_2}) = 0$ for $\sigma_1 \neq \sigma_2^*$;
- $(B_{\sigma_1}, B_{\sigma_2}, B_{\sigma_3}) = 0$ for $\Omega(\sigma_1, \sigma_2, \sigma_3) = \emptyset$
- $\sum_{i,j} (x_1, x_2, b_i^{(1,2|3,4)}) F^{ij}_{(1,2|3,4)} (b_j^{(3,4|1,2)}, x_3, x_4) = \sum_{i,j} (x_4, x_1, b_i^{(4,1|2,3)}) F^{ij}_{(4,1|2,3)} (b_j^{(2,3|4,1)}, x_2, x_3)$.

Here $x_k \in B_{\sigma_k}$, $\{b_i^{(s,t|k,r)}\}$ is a basis of $B_{\sigma_{(s,t|k,r)}}$ and $F^{ij}_{(s,t|k,r)}$ is the inverse matrix for $(b_i^{(s,t|k,r)}, b_j^{(k,r|s,t)})$.

We will consider B_* as an algebra with the multiplication $(x_1 x_2, x_3) = (x_1, x_2, x_3)$, for $x_k \in B_{\sigma_k}$. The axiom $\sum_{i,j} (x_1, x_2, b_i^{(1,2|3,4)}) F^{ij}_{(1,2|3,4)} (b_j^{(3,4|1,2)}, x_3, x_4) = \sum_{i,j} (x_4, x_1, b_i^{(4,1|2,3)}) F^{ij}_{(4,1|2,3)} (b_j^{(2,3|4,1)}, x_2, x_3)$ is equivalent to associativity for the algebra B_*. Moreover B_* is a Frobenius Algebra in the sense of [7] if its dimension is finite.

3.2. **Topology-algebra correspondence.**

Theorem 3.1. *Let* $\mathcal{F} = \{\Phi_{\Omega} : V_{\Omega} \to \mathbb{K}\}$ *be a Brane Topological Field Theory on a tuple of finite-dimensional vector spaces* $\{B_{\sigma} | \sigma \in \Sigma\}$*. Then the poli-linear forms*

- $(x', x'') = \Phi_{\Omega(\sigma_1, \sigma_2)}(x'_{q_1} \otimes x''_{q_2})$*, where* $x' \in B_{\sigma_1}$*,* $x'' \in B_{\sigma_2}$
- $(x', x'', x''') = \Phi_{\Omega(\sigma_1, \sigma_2, \sigma_3)}(x'_{q_1} \otimes x''_{q_2} \otimes x'''_{q_3})$*, где* $x' \in B_{\sigma_1}$*,* $x'' \in B_{\sigma_2}$*,* $x''' \in B_{\sigma_3}$*.*

generate a structure of Colored Frobenius Algebra on $B_{\star} = \bigoplus_{\sigma \in \Sigma} B_{\sigma}$*.*

Proof. Only the last axiom is not obvious. Let us consider a brane complex $\Omega \in \Omega(\sigma_1, \sigma_2, \sigma_3, \sigma_4)$, and a cut between the pairs of vertexes σ_1, σ_2 and σ_3, σ_4. Then the cut-invariant axiom and lemma 2.2 give

$\sum_{i,j}(x_1, x_2, b_i^{(1,2|3,4)})F^{ij}_{(1,2|3,4)}(b_j^{(3,4|1,2)}, x_3, x_4) = \Phi_\Omega(x_1, x_2, x_3, x_4)$. Similarly, $\sum_{i,j}(x_4, x_1, b_i^{(4,1|2,3)})F^{ij}_{(4,1|2,3)}(b_j^{(2,3|4,1)}, x_2, x_3) = \Phi_\Omega(x_1, x_2, x_3, x_4)$
□

Theorem 3.2. *Let $B_\star = \bigoplus_{\sigma\in\Sigma} B_\sigma$ be a Colored Frobenius Algebra with poli-linear forms $(.,.)$ and $(.,.,.)$. Then it generates a Brane Topological Field Theory on $\{B_\sigma|\sigma \in \Sigma\}$ by means of following construction. Fix a basis $\{b_i^\sigma\}$ of any vector space B_σ, $\sigma \in \Sigma$. Consider the matrix F_σ^{ij} that is the inverse matrix for $F_{ij}^\sigma = (b_i^\sigma, b_j^{\sigma^*})$. Define the linear functionals for any connected brane complexes by*

- $\Phi_{\Omega(\sigma_1,\sigma_2\dots,\sigma_n)}(x^1_{q_1} \otimes x^2_{q_2} \otimes \dots \otimes x^n_{q_n}) = \sum_{\varsigma_1,\varsigma_2,\dots,\varsigma_{n-3}\in\Sigma}(x^1, x^2, b_{i_1}^{\varsigma_1})F^{\varsigma_1}_{i_1 j_1}\ (b_{j_1}^{\varsigma_1^*}, x^3_{q_3}, b_{i_2}^{\varsigma_2})$ $F^{\varsigma_2}_{i_2 j_2}(b_{j_2}^{\varsigma_2^*}, x^4_{q_4}, b_{i_3}^{\varsigma_3})\dots\dots F^{\varsigma_{n-4}}_{i_{n-4} j_{n-4}}(b_{j_{n-4}}^{\varsigma_{n-4}^*}, x^{n-2}_{q_{n-2}}, b_{i_{n-3}}^{\varsigma_{n-3}})F^{\varsigma_{n-3}}_{i_{n-3} j_{n-3}}(b_{j_{n-3}}^{\varsigma_{n-3}^*}, x^{n-1}_{q_{n-1}}, x^n_{q_n})$, *where $x^i_{q_i} \in B_{\sigma_i}$.*

Then the family $\{\Phi_{\Omega(\sigma_1,\sigma_2\dots,\sigma_n)}\}$ generates a Brane Topological Field Theory.

Proof. The topological invariance follows from the invariance under cyclic renumbering the vertices of Ω. The invariance under the renumbering $q_i \mapsto q_j$ $j \equiv i + 1(\mathrm{mod}2)$ follows from the last axiom for the tree-linear form. The cut invariance follows directly from the definition of Φ if we renumber the vertices marking the cut divide the vertices $q_1, q_2, \dots q_k$ and $q_{k+1}, q_{k+2}, \dots q_n$.
□

These two theorems determine the one-to-one correspondence between Brane Topological Field Theories and isomorphic classes of Colored Frobenius Algebras.

4. HURWITZ NUMBERS OF COLORED COMPLEXES

4.1. **Hurwitz numbers.** Classical Hurwitz numbers are weighted number of brunched coverings over closed surface with prescribed types of critical values [8]. Let us define some analog of Hurwitz numbers for brane complexes.

A brunched covering between simple CW-complexes $f : \widetilde{\Omega} \to \Omega$ is called *simple covering of degree d* or *simple d-covering* it is a local homeomorphism on any cell, preimage of any cell consists of unions of cells, and the number of cells from preimage is d for any cell of higher dimension. Simple coverings $f' : \widetilde{\Omega}' \to \Omega$ and $f'' : \widetilde{\Omega}'' \to \Omega$ are called *isomorphic* if there exists an isomorphism $\widetilde{\varphi} : \widetilde{\Omega}' \to \widetilde{\Omega}''$ such that $f' = f''\widetilde{\varphi}$. Denote by $\mathsf{Aut}(f)$ the automorphisms group of a simple covering f.

Let Ω' and Ω'' colored complexes. Simple coverings $f' : \widetilde{\Omega}' \to \Omega'$ and $f'' : \widetilde{\Omega}'' \to \Omega''$ are called *equivalent* if there exists isomorphisms $\widetilde{\varphi} : \widetilde{\Omega}' \to \widetilde{\Omega}''$ and $\varphi : \Omega' \to \Omega''$ such that $\varphi f' = f''\widetilde{\varphi}$. Denote by Υ^d_Ω the set of equivalent classes of simple d-coverings over a colored complex Ω.

A simple d-covering $f : \widetilde{\Omega} \to \Omega$ over a brane complex Ω generate a simple d-covering $f_q : \widetilde{\Omega}_q \to \sigma_q$ over the vertex complex σ_q of a vertex q. The equivalence class of f_q is called a *local invariant* of f at q. Consider a set $\{\beta_q\} = \{\beta_q \in \Upsilon^d_{\sigma_q} | q \in \Omega_b\}$. Denote by $\Upsilon^d_\Omega(\{\beta_q\})$ the set of isomorphic classes of simple d-coverings $f : \widetilde{\Omega} \to \Omega$ with local invariants $f_q = \beta_q$ for all vertexes $q \in \Omega_b$. The weighted number of simple d-covering

$$H^d(\Omega, \{\beta_q\}) = \sum_{f\in\Upsilon^d_\Omega(\{\beta_q\})} 1/|\,\mathsf{Aut}(f)|$$

is called *Hurwitz number of simple d-coverings*

4.2. **Hurwitz Brane Topological Field Theory.** For any $\sigma \in \Sigma = \Sigma(D, S)$, denote by C^d_σ the vector space generated by the set Υ^d_σ. The involution $\sigma \mapsto \sigma^*$ generates the involution $C^d_\sigma \mapsto C^d_{\sigma^*}$. Hurwitz numbers generate a family of linear functions $\mathcal{H}^d = \{\Phi_\Omega : V_\Omega \to \mathbb{K}\}$ on $\{V_\Omega = (\otimes_{q \in \Omega_b} C^d_q)\}$, depending from brane complexes Ω.

Theorem 4.1. *$\mathcal{H}^d$ is a Brane Topological Field Theory.*

Proof. It is not evident only the cut-invariant axiom. Fix a cut $\sigma \subset \Omega \in \Omega(\sigma_1, ..., \sigma_m)$ and a covering $\beta \in \Upsilon^d_\sigma$. Let $\Omega \setminus \sigma = \Omega_+ \cup \Omega_-$. Put $\overline{\Omega}_+ = \Omega_+ \cup \sigma_+ \in \Omega(\sigma_1, ..., \sigma_k, \sigma)$ and $\overline{\Omega}_- = \Omega_+ \cup \sigma_+ \in \Omega(\sigma, \sigma_{k+1}, ..., \sigma_m)$. The simple covering $f : \widetilde{\Omega} \to \Omega$ generates the simple coverings $f_\pm : (\widetilde{\Omega}_\pm \cup f^{-1}(\sigma)_\pm)) \to \overline{\Omega}_\pm$, where $\widetilde{\Omega}_\pm = f^{-1}(\Omega_\pm)$. Consider the set $M^\beta_\Omega(\beta_1, ..., \beta_m)$ of simple coverings $f : \widetilde{\Omega} \to \Omega$, such that local invariants of f at vertexes of Ω are β_i and the covering $f_+|_{f^{-1}(\sigma)_+}$ is equivalent to β.

The coverings from $M^\beta_\Omega(\beta_1, ..., \beta_m)$ are coverings that appear from coverings $f_+ : (\widetilde{\Omega}_+ \cup f^{-1}(\sigma)_+)) \to \overline{\Omega}_+$ and $f_- : (\widetilde{\Omega}_- \cup f^{-1}(\sigma)_-)) \to \overline{\Omega}_-$ by gluing according to an equivalence $f_+|_{f^{-1}(\sigma)_+} \to f_-|_{f^{-1}(\sigma)_-}$. Moreover, different equivalences $f_+|_{f^{-1}(\sigma)_+} \to f_-|_{f^{-1}(\sigma)_-}$ generate isomorphic coverings if one of it appear from other by automorphisms of f_+ and f_-. Thus, $\sum_{f \in M^\beta_\Omega(\beta_1,...,\beta_m)} \frac{1}{|\mathsf{Aut}(f)|} = \sum_{f_+ \in \Upsilon^d_{\overline{\Omega}_+}(\beta_1,...,\beta_k,\beta), f_- \in \Upsilon^d_{\overline{\Omega}_-}(\beta,\beta_{k+1},...,\beta_m)} \frac{|\mathsf{Aut}(\beta)|}{|\mathsf{Aut}(f_+)||\mathsf{Aut}(f_-)|} = \sum_{f_+ \in \Upsilon^d_{\overline{\Omega}_+}(\beta_1,...,\beta_k,\beta)} \frac{1}{|\mathsf{Aut}(f_+)|} |\mathsf{Aut}(\beta)| \sum_{f_- \in \Upsilon^d_{\overline{\Omega}_-}(\beta,\beta_{k+1},...,\beta_m)} \frac{1}{|\mathsf{Aut}(f_-)|}$. Therefore $H^d(\Omega, \{\beta_1, ...\beta_m\}) = \sum_{\beta \in \Upsilon^d_\sigma} \sum_{f \in M^\beta_\Omega|(\beta_1,...,\beta_m)} \frac{1}{|\mathsf{Aut}(f)|} = \sum_{\beta \in \Upsilon^d_\beta} H(\Omega_+, \{\beta_1, ...\beta_k, \beta\}) |\mathsf{Aut}(\beta)| H(\Omega_-, \{\beta, \beta_{k+1}, ...\beta_m\})$. Moreover $H(\Omega, \{\beta, \gamma\}) = \frac{\delta_{\beta,\gamma^*}}{|\mathsf{Aut}(\beta)|}$.
□

It is follow from theorem 3.1 that the theory $\mathcal{H}^d$ generates a Colored Frobenius Algebra C^d_*. It defines a multiplication between isomorphic classes of simple d-coverings of colored complexes. Theorem 3.2 gives a formula for calculation of Hurwitz numbers of simple coverings in terms of these algebra. The algebra C^d_* is an hight dimensional analog of the algebra bi-party graphs from [2, 3]. Algebra C^1_* is an algebra of brane complexes.

5. GROUP FOAM

5.1. **Hurwitz numbers of G-coverings.** Fix a finite group G. A homomorphism $G \to \mathsf{Aut}(\widetilde{\Omega})$ to the group of automorphism of simple CW-complex $\widetilde{\Omega}$ is called a *G-action* on $\widetilde{\Omega}$, if the set of fixed points of any $g \in G$ is sum of some full cell of $\widetilde{\Omega}$. For brevity we shall identify elements of G and its images in $\mathsf{Aut}(\widetilde{\Omega})$.

We say that a map $f : \widetilde{\Omega} \to \Omega$ between CW-complexes is *G-covering* if $f = f_* r$, where $r : \widetilde{\Omega} \to \widetilde{\Omega}/G$ is the natural projection by some G-action and $f_* : \widetilde{\Omega}/G \to \Omega$ is an isomorphism. G-coverings $f' : \widetilde{\Omega}' \to \Omega$ and $f” : \widetilde{\Omega}” \to \Omega$ are called *isomorphic*, if there exists an isomorphism $\widetilde{\varphi} : \widetilde{\Omega}' \to \widetilde{\Omega}”$ such that $\widetilde{\varphi} g = g \widetilde{\varphi}$, for $g \in G$, and $f' = f” \widetilde{\varphi}$. Denote by $\mathsf{Aut}^G(f)$ the automorphisms group of a G-covering f.

Let Ω' and $\Omega”$ colored complexes. G-coverings $f' : \widetilde{\Omega}' \to \Omega'$ and $f” : \widetilde{\Omega}” \to \Omega”$ are called *equivalent*, if there exists isomorphisms $\widetilde{\varphi} : \widetilde{\Omega}' \to \widetilde{\Omega}”$ and $\varphi : \Omega' \to \Omega”$ such that $\widetilde{\varphi} g = g \widetilde{\varphi}$, for $g \in G$, and $\varphi f' = f” \widetilde{\varphi}$. Denote by Υ^G_Ω the set of equivalent classes of G-coverings over a colored complex Ω.

A G-covering $f : \widetilde{\Omega} \to \Omega$ over a brane complex Ω generates a G-covering $f_q : \widetilde{\Omega}_q \to \sigma_q$ over the vertex complex σ_q of a vertex q. The equivalence class of f_q is called a *local invariant* of f at q. Consider a set $\{\beta_q\} = \{\beta_q \in \Upsilon^G_{\sigma_q} | q \in \Omega_b\}$. Denote by $\Upsilon^G_\Omega(\{\beta_q\})$ the set of isomorphic classes of G-coverings $f : \widetilde{\Omega} \to \Omega$ with local invariants $f_q = \beta_q$ for all vertexes $q \in \Omega_b$. The weighted number of G-covering

$$H^G(\Omega, \{\beta_q\}) = \sum_{f \in \Upsilon^G_\Omega(\{\beta_q\})} 1/|\operatorname{\mathsf{Aut}}^G(f)|$$

is called *Hurwitz Number of G-coverings*

5.2. **Brane Topological Field Theory of G-coverings.** For any $\sigma \in \Sigma = \Sigma(D, S)$, denote by C^G_σ the vector space generated by the set Υ^G_σ. The involution $\sigma \mapsto \sigma^*$ generates the involution $C^G_\sigma \mapsto C^G_{\sigma^*}$. Hurwitz numbers generate a family of linear functions $\mathcal{H}^G = \{\Phi_\Omega : V_\Omega \to \mathbb{K}\}$ on $\{V_\Omega = (\otimes_{q \in \Omega_b} C^G_q)\}$, depending from brane complexes Ω. By repeating the proof of theorem 4.1 we obtain

Theorem 5.1. *$\mathcal{H}^G$ is a Brane Topological Field Theory.*

It is follow from theorem 5.1 that $\mathcal{H}^G$ generates a Colored Frobenius Algebra C^G_* (*G-foam*). It is obviously that its isomorphic class depends only on G and $|S|$. Thus, G-foams are invariants of G that describe the family of subgroups of G. The restriction of G-foam on the category of 2-dimensional disks (see remark 2.1) were consider in [3]. The algebra C^G_* coincides with the algebra C^d_* from previous section, if G is the symmetrical group S_d.

REFERENCES

[1] Alexeevski A., Natanzon S., Noncommutative two-dimensional topological field theories and Hurwitz numbers for real algebraic curves. Selecta Math., New ser. v.12,n.3, 2006, p. 307-377 (arXiv: math.GT/0202164).

[2] Alexeevski A., Natanzon S., Algebra of Hurwitz numbers for seamed surfaces, Russian Math.Surveys, 61 (4) (2006), 767-769

[3] Alexeevski A., Natanzon S., Algebra of bipartite graphs and Hurwitz numbers of seamed surfaces. Math.Russian Izvestiya RAN: Ser. Mat. 72:4 627-643.

[4] Alexeevski A., Natanzon S., Hurwitz numbers for regular coverings of surfaces by seamed surfaces and Cardy-Frobenius algebras of finite groups, Amer. Math. Soc. Transl. (2) Vol 224, (2008), 1-25 (arXiv: math/07093601)

[5] Atiyah M., Topological Quantum Field Theories, Inst. Hautes Etudes Sci. Publ. Math., 68 (1988), 175-186.

[6] Bucstaber V, Rees E., Frobenius n-homomorphisms, transfers and branched coverings, arXiv: math/0608120

[7] Faith C., Algebra II Ring theory, Springer-Verlag, 1976

[8] Hurwitz A., Über Riemann'sche Flächen mit gegeben Verzweigungspunkten, Math., Ann., Bn.39 (1891), 1-61.

[9] Lazaroiu C.L., On the structure of open-closed topological field theory in two-dimensions, Nucl.Phys. B 603, 2001, 497-530.

[10] Moore G., Segal G, D-branes and K-theory in 2D topological field theory, arXiv:hep-th/0609042 (2006)

[11] Natanzon S., Cyclic foam topological fied theory. arXiv:0712.3557

[12] Witten.E, Quantum Field theory. Commun.Math.Phys.117(1988),353-386